Completeness and Termination of Tableau Calculus for Undirected Graphs

Yuki Nishimura^[0009-0000-0475-1168] and Tsubasa Takagi^[0000-0001-9890-1015]

School of Computing, Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro-ku, Tokyo 152-8550 Japan {nishimura.y.as,takagi.t.ah}@m.titech.ac.jp

Abstract. Hybrid logic is a modal logic with additional operators specifying nominals and is highly expressive. For example, there is no formula corresponding to the irreflexivity of Kripke frames in basic modal logic, but there is in hybrid logic. Irreflexivity is significant in that irreflexive and symmetric Kripke frames can be regarded as undirected graphs reviewed from a graph theoretic point of view. Thus, the study of the hybrid logic with axioms corresponding to irreflexivity and symmetry can help to elucidate the logical properties of undirected graphs. In this paper, we formulate the tableau method of the hybrid logic for undirected graphs. Our main result is to show the completeness theorem and the termination property of the tableau method, which leads us to prove the decidability.

Keywords: Hybrid logic \cdot Tableau calculus \cdot Termination \cdot Undirected graph \cdot Orthoframe.

1 Introduction

Hybrid logic is an extension of basic modal logic with additional propositional symbols called nominals and an operator called a satisfaction operator @. A nominal is a formula that is true only in a single possible world in a Kripke model. Nominals are regarded as syntactic names of possible worlds. Hybrid logic is highly expressive because nominals allow arbitrary possible worlds to be designated. Hybrid logic was first invented by Prior [17, 18] and has developed in various ways since then (see [1, 5, 9, 10, 15]).

One of the advantages of hybrid logic compared to basic modal logic is that we can treat the irreflexivity of Kripke frames. In basic modal logic, no axiom corresponds to the irreflexivity of the model [12]. In hybrid logic, however, we have a simple axiom, $@_i \neg \Diamond i$, where *i* denotes a nominal [6].

This advantage makes hybrid logic well-suited for dealing with undirected graphs, namely relational structures with irreflexivity and symmetry. There are some logics whose semantics are given by undirected graphs; for example, orthoframes used in (the minimal) quantum logic [13] and social networks used in Facebook logic [20, 21] can be regarded as undirected graphs.

The main result of this paper is to propose a tableau calculus that is sound and complete with respect to the class of irreflexive and symmetric frames. Moreover, we show that any tableau can be constructed within a finite time, called termination property. For a given logic, if its tableau calculus is terminating, we can derive that it is decidable. That is, whether an arbitrary formula is valid or not is automatically shown. In addition, if the formula is not valid, we can construct a countermodel from the tableau to refute the validity of the formula.

We proceed as follows: Section 2 introduces the semantics of hybrid logic, and Section 3 shows the tableau system of basic hybrid logic. After that, we introduce the tableau calculus corresponding to undirected graphs in Section 4.

Related Work

Tableau Calculi for Hybrid Logic. Bolander and Blackburn [7,8] studied tableau calculi for hybrid logic. Bolander & Blackburn (2009) [8] proposed a tableau calculus for hybrid logic based on $\mathbf{K4}$, and claimed that the loop-checking approach can be extended in it to the hybrid logics based on $\mathbf{S4}$ and $\mathbf{S5}$. This study, however, addressed a combination (namely, the combination of (irr) and (sym)) that they have not conducted because they considered them challenging: in [8], they wrote, "But then we are faced with the task of combining such conditions as (irr), (sym), (asym), (antisym), (intrans), (uniq) and (tree) with (trans), and here matters are likely to be much trickier." As we have already mentioned, we have paid particular attention to the combination of (irr) and (sym) because it characterizes undirected graphs as Kripke frames and is, therefore, significant from a graph-theoretic point of view.

Hybrid Logic for Graph Theory. Examples of research on graph theory using hybrid logic can be found in [2, 3, 11]. They studied hybrid graph logic, a hybrid logic with a special operator \diamond^+ , as a tool for analyzing some properties of graph theory. In particular, for undirected graphs, [16] discussed how to express the planarity of undirected graphs (note that [16] also showed the decidability of hybrid logic for undirected graphs). All of these studies adopted Hilbert-style axiomatization as their proof system. On the other hand, we adopt tableau calculus as the proof system. As is well known, tableau calculi have an advantage compared to Hilbert-style axiomatizations in that the proof construction procedure is automatic.

Bulldozing Method. Blackburn (1990) [4] used the bulldozing method in proving the completeness of hybrid logic with the axiom corresponding to irreflexivity. Our work shows that the bulldozing method also works in dealing with the tableau calculus for hybrid logic with a rule of irreflexivity. The work of Blackburn (1990) [4] showed that none of the reflexive points in the model created by a maximal consistent set has a nominal and checked that there is no matter to apply bulldozing. In our work, however, all the models created by an open saturated branch are named — each world makes at least one nominal true. We show that this situation causes no problem since all the nominals true in reflexive worlds are not an element that determines the validity of a formula but just "labels" to point to worlds. In this sense, our work is an extension of [4].

2 Kripke Semantics of Hybrid Logic

Here, we review a hybrid logic with an operator @ (see [9] for more details.)

Definition 1 (language). We have a countable infinite set **Prop** of propositional variables¹ and another countable infinite set **Nom** of nominals which is disjoint from **Prop**. The set of formulas of hybrid logic is defined inductively as follows:

$$\varphi ::= p \mid i \mid \neg \varphi \mid \varphi \land \varphi \mid \Diamond \varphi \mid @_i \varphi,$$

where $p \in \mathbf{Prop}$ and $i \in \mathbf{Nom}$.

We use the following abbreviations:

$$\varphi \lor \psi \coloneqq \neg (\neg \varphi \land \neg \psi), \ \varphi \to \psi \coloneqq \neg (\varphi \land \neg \psi), \ \Box \varphi \coloneqq \neg \Diamond \neg \varphi.$$

Definition 2. A Kripke model (we call it model, in short) $\mathcal{M} = (W, R, V)$ is defined as follows:

- -W is a non-empty set.
- R is a binary relation on W. That is, $R \subseteq W \times W$.
- V is a function $V : \operatorname{Prop} \cup \operatorname{Nom} \to \mathcal{P}(W)$ such that $V(i) = \{w\}$ for some $w \in W$ for each $i \in \operatorname{Nom}$, where $\mathcal{P}(W)$ denotes the powerset of W.

Furthermore, we call $\mathcal{F} = (W, R)$ that satisfies the conditions above a Kripke frame (or shortly frame).

This definition reflects the property of nominals: one nominal is true in only one world. In this paper, we write wRv to mean $(w, v) \in R$. Moreover, We abbreviate $w \in W$ such that $V(i) = \{w\}$ by i^V .

Definition 3. Given a model \mathcal{M} , a possible world w in \mathcal{M} , and a formula φ , the satisfaction relation $\mathcal{M}, w \models \varphi$ is defined inductively as follows:

 $\begin{array}{ll} \mathcal{M}, w \models p & i\!f\!f \quad w \in V(p), where \ p \in \mathbf{Prop}; \\ \mathcal{M}, w \models i & i\!f\!f \quad w = i^V, where \ i \in \mathbf{Nom}; \\ \mathcal{M}, w \models \neg \varphi & i\!f\!f \quad not \ \mathcal{M}, w \models \varphi; \\ \mathcal{M}, w \models \varphi \wedge \psi & i\!f\!f \quad \mathcal{M}, w \models \varphi \ and \ \mathcal{M}, w \models \psi; \\ \mathcal{M}, w \models \Diamond \varphi & i\!f\!f \quad there \ is \ v \ such \ that \ wRv \ and \ \mathcal{M}, v \models \varphi; \\ \mathcal{M}, w \models @_i \varphi & i\!f\!f \quad \mathcal{M}, i^V \models \varphi. \end{array}$

Definition 4. A formula φ is valid on \mathcal{M} , denoted by $\mathcal{M} \models \varphi$, if $\mathcal{M}, w \models \varphi$ holds for all $w \in W$. Also, if $\mathcal{M}, w \models \varphi$ holds for all models \mathcal{M} and all of its worlds w, we say that φ is valid, and we write $\models \varphi$.

¹ We suppose that **Prop** is countable, following [9].

$$\frac{\underline{@}_{i}\neg j}{\underline{@}_{j}j} \ [\neg] \qquad \frac{\underline{@}_{i}\neg \neg \varphi}{\underline{@}_{i}\varphi} \ [\neg\neg] \qquad \frac{\underline{@}_{i}(\varphi \wedge \psi)}{\underline{@}_{i}\psi} \ [\wedge] \qquad \frac{\underline{@}_{i}\neg (\varphi \wedge \psi)}{\underline{@}_{i}\neg \varphi \mid \underline{@}_{i}\neg \psi} \ [\neg\wedge]$$

*1: $j \in \mathbf{Nom}$ does not occur in the branch.

*2: This rule can be applied only one time per formula.

*3: The formula above the line is not an accessibility formula. Here, an *accessibility* formula is the formula of the form $@_i \Diamond j$ generated by $[\Diamond]$, where j is a new formula.

In these rules, the formulas above the line show the formulas that have already occurred in the branch, and the formulas below the line show the formulas that will be added to the branch. The vertical line in the $[\neg \land]$ rule means that the branch splits to the left and right.

Fig. 1: The rules of TAB

3 Internalized Tableau Calculus for K(@)

 $\mathbf{K}(@)$ is an axiomatization of hybrid logic with an operator @, based on the simplest normal modal logic \mathbf{K} (see [5], for example.) We write **TAB** to indicate the tableau calculus for hybrid logic $\mathbf{K}(@)$. This section is based on [7].

Definition 5. A tableau is a well-founded tree constructed in the following way:

- Start with a formula of the form $@_i \varphi$ (called the root formula), where φ is a formula of hybrid logic and $i \in \mathbf{Nom}$ does not occur in φ .
- For each branch, extend it by applying rules (see Definition 7) to all nodes as often as possible. However, we can no longer add any formula in a branch if at least one of the following conditions is satisfied:
 - *i)* Every new formula generated by applying any rule already exists in the branch.
 - *ii)* The branch is closed (see Definition 6.)

Here, a branch means a maximal path of a tableau. If a formula φ occurs in a branch Θ , we write $\varphi \in \Theta$.

Definition 6. A branch of a tableau Θ is closed if there is some φ and $i \in \mathbf{Nom}$ such that $@_i \varphi, @_i \neg \varphi \in \Theta$. We say that Θ is open if it is not closed. A tableau is called closed if all branches in the tableau are closed.

Definition 7. We provide the rules of **TAB** in Figure 1.

Definition 8 (provability). Given a formula φ , we say that φ is provable in **TAB** if there is a closed tableau whose root formula is $@_i \neg \varphi$, where $i \in \mathbf{Nom}$ does not occur in φ .

[7] shows two significant properties of **TAB**. One is the termination property, and the other is completeness.

Theorem 1 (termination). TAB has the termination property. That is, for every tableau in **TAB**, all branches of it are finite.

Theorem 2 (completeness). **TAB** is complete for the class of all frames.

4 Tableau Calculus for IB(@)

In this paper, we discuss the hybrid logic corresponding to irreflexive and symmetric models. We call the models orthomodels, following [13].

Definition 9. An orthoframe is a Kripke frame $\mathcal{F} = (W, R)$, where R is irreflexive and symmetric. Also, an orthomodel is $\mathcal{M} = (\mathcal{F}, V)$, where \mathcal{F} is an orthoframe.

IB(@) is a hybrid logic corresponding to the class of all orthoframes. We can construct it by adding the following two axioms to the hybrid logic K(@):

4.1 Constructing Tableau Calculus

Our aim is to construct a tableau calculus corresponding to IB(@), keeping termination. First, we add the following rule that reflects the symmetry:

$$\frac{@_i \Box \varphi}{@_j \diamondsuit i} \\ \frac{@_j \diamondsuit i}{@_j \varphi} \ [\Box_{sym}]$$

However, this rule allows us to build an infinite branch.

Example 1. Consider an example of Figure 2 (formulas with * are accessibility formulas). In this branch, nominals are generated infinitely as i_1, i_3, i_5, \ldots and i_2, i_4, i_6, \ldots

In this case, the nominals i_1, i_3, i_5, \ldots play the same role. For any nominal i_{2n+1} $(n \ge 0)$ occurring in Θ , we have $@_{i_{2n+1}} \diamondsuit \diamondsuit p, @_{i_{2n+1}} \diamondsuit p \in \Theta$. Then, we have to prohibit the creation of such redundant nominals.

The solution to this problem has already been proposed in [7, Section 5.2] and [8, Section 7]. We need to define some concepts and show their properties to introduce the restriction.

First, we define the notion of quasi-urfather. In a sense, it can be regarded as a representative of nominals.

$1.@_{i_0}(\diamondsuit \Diamond i \land @_i \Box \diamondsuit \Diamond i)$	
$2.@_{i_0} \diamondsuit \diamond i$	(1)
$3.@_{i_0}@_i\Box\Diamond\Diamond i$	(1)
$4.@_{i_0} \diamondsuit i_1^*$	(2)
$5.@_{i_1} \diamondsuit i$	(2)
$6.@_i \Box \diamondsuit \diamondsuit i$	(3)
$7.@_{i_1} \diamondsuit {i_2}^*$	(5)
$8.@_{i_2}i$	(5)
$9.@_{i_1} \diamondsuit \diamond i$	$(5, 6, [\Box_{sym}])$
$10.@_{i}i$	(8)
$11.@_{i_1} \diamondsuit {i_3}^*$	(9)
$12.@_{i_3} \diamondsuit i$	(9)
$13.@_{i_3} \diamondsuit i_4^*$	(12)
$14.@_{i_4}i$	(12)
$15.@_{i_3} \diamondsuit \diamond i$	$(6, 12, [\Box_{sym}])$
$16.@_{i_3} \diamondsuit {i_5}^*$	(15)
$17.@_{i_5} \diamondsuit i$	(15)
$18.@_{i_5} \diamondsuit i_6^*$	(17)
$19.@_{i_6}i$	(17)
$20.@_{i_5} \diamondsuit \diamond i$	$(6, 17, [\Box_{sym}])$
:	
•	

Fig. 2: Non-terminating tableau of **TAB** with only $[\Box_{sym}]$

Definition 10 (quasi-subformula). Given two formulas of the form $@_i \varphi$ and $@_i \psi$, $@_i \varphi$ is a quasi-subformula of $@_i \psi$ if one of the following conditions holds:

- $-\varphi$ is a subformula of ψ .
- $-\varphi$ has the form $\neg \chi$, and χ is a subformula of ψ .

Lemma 1. Let \mathcal{T} be a tableau. For every formula of the form $@_i \varphi$ in \mathcal{T} , it is either a quasi-subformula of the root formula of \mathcal{T} or an accessibility formula.

Proof. See [7, Lemma 6.1].

Definition 11. Let Θ be a branch of a tableau. For every nominal occurring in Θ , the set of true formulas at *i*, denoted by $T^{\Theta}(i)$ is defined as follows:

 $T^{\Theta}(i) = \{ \varphi \mid @_i \varphi \in \Theta \text{ and } @_i \varphi \text{ is a quasi-subformula of a root formula} \}.$

Completeness and Termination of Tableau Calculus for Undirected Graphs

Second, we define a generating relation on nominals.

Definition 12. Let Θ be a branch of tableau and let i and j be nominals occurring in Θ . If j is introduced by applying $[\diamondsuit]$ to a formula $@_i \diamondsuit \varphi$, we say that j is a generated nominal from i (denotation: $i \prec_{\Theta} j$).

Note that the only rule generating a new nominal is $[\diamondsuit]$. Since all the accessibility formulas emerge in a branch if and only if $[\diamondsuit]$ is applied to some formula in the branch, $i \prec_{\Theta} j$ is equivalent to there being an accessibility formula $@_i \diamondsuit j$ in Θ .

Definition 13 (quasi-urfather, [7, Definition 5.7]). Let Θ be a branch of a tableau. We call nominals i, j twins in Θ if $T^{\Theta}(i) = T^{\Theta}(j)$. Also, we call a nominal *i* quasi-urfather on Θ if there are no twins j, k such that $j \neq k$ and $j, k \prec_{\Theta}^{*} i$. Here, \prec_{Θ}^{*} is a reflexive and transitive closure of \prec_{Θ} .

Then, we can declare the restriction, which keeps the termination property of our tableau calculus.

 (\mathcal{D}) The rule $[\diamondsuit]$ can only be applied to a formula $@_i \varphi$ on a branch Θ if i is a quasi-urfather.

Example 2. With (\mathcal{D}) , the tableau in Figure 2 stops before adding 16. $@_{i_3} \diamond_{i_5}$. In this branch, up to 15. $@_{i_3} \diamond \diamond_i$, nominals i_1 and i_3 are twins, since $T_{\Theta}(i_1) = T_{\Theta}(i_3) = \{ \diamond_i, \diamond \diamond_i \}$. Then, i_3 is not a quasi-urfather, so we can no longer apply $[\diamond]$ to 15. $@_{i_3} \diamond \diamond_i$.

The next thing to do is the addition of rules that address irreflexivity. The solution, which is quite similar to that of [8], is straightforward: we add the form of $@_i \neg \Diamond i$ to the branch for every nominal i.

 (\mathcal{I}) For any nominal *i* occurring in Θ , we add a formula $@_i \neg \Diamond i$.

Example 3. Consider an example of Figure 3. The branch with the root formula $@_i \diamond (i \land p)$ is closed owing to (\mathcal{I}) .

Now, we are ready to construct the tableau calculus for IB(@).

Definition 14. $\operatorname{TAB}_{\operatorname{IB}(@)}$ (or simply $\operatorname{TAB}_{\operatorname{IB}}$) is the tableau calculus made by adding $[\Box_{sym}], (\mathcal{D}), and (\mathcal{I})$ to TAB . Moreover, if there is a closed tableau in $\operatorname{TAB}_{\operatorname{IB}}$ whose root formula is $@_i \neg \varphi$ where $i \in \operatorname{Nom}$ does not occur in φ , we say that φ is provable in $\operatorname{TAB}_{\operatorname{IB}}$.

4.2 Termination and Completeness

First, we show the termination property.

Theorem 3. TAB_{IB} has the termination property.

$1.@_i \diamondsuit (i \land p)$	
$2.@_i \neg \diamondsuit i$	(\mathcal{I})
$3.@_i \diamondsuit j^*$	(1)
$4.@_j(i \wedge p)$	(1)
$5.@_j \neg \diamondsuit j$	(\mathcal{I})
$6.@_j \neg i$	(2, 3)
$7.@_ji$	(4)
$8.@_jp$	(4)
×	

Fig. 3: A closed tableau owing to (\mathcal{I})

Proof. Suppose there is an infinite branch Θ . Then, there is an infinite row of nominals as follows (for more detail, see the proof of [7, Lemma 6.5]):

$$i_0 \prec_{\Theta} i_1 \prec_{\Theta} i_2 \prec_{\Theta} \cdots$$

For Θ and its root formula $@_i \varphi$, define Q and n as follows:

$$Q = \{ \psi \mid @_i \psi \text{ is a quasi-subformula of } @_i \varphi \},$$

$$n = |Q|.$$

Also, let Θ' be a fragment of Θ up to, but not including, the first occurrence of i_{2^n+1} . Then i_{2^n+1} is generated by applying $[\diamondsuit]$ to some $@_{i_{2^n}} \diamondsuit \varphi$. Taking (\mathcal{D}) into consideration, i_{2^n} is a quasi-urfather. However, since all of $T^{\Theta'}(i_0), T^{\Theta'}(i_1), \ldots, T^{\Theta'}(i_{2^n})$ are subsets of Q and n is

However, since all of $T^{\Theta'}(i_0), T^{\Theta'}(i_1), \ldots, T^{\Theta'}(i_{2^n})$ are subsets of Q and n is the cardinality of Q, there is a pair $1 \leq l, m \leq 2^n$ such that $T^{\Theta'}(i_l) = T^{\Theta'}(i_m)$ (pigeonhole principle). Therefore, we find the twins i_l and i_m such that $i_l, i_m \prec_{\Theta}^*$ i_{2^n} , but that contradicts that i_{2^n} is a quasi-urfather.

Next, we prove completeness. The basic strategy is the same as in [7, Section 5.2]. That is, define an *identity urfather* (see [7, Definition 5.10]) and use it to create a model from an open saturated branch of tableau (when constructing the model, make sure that the relation R^{Θ} is symmetric). Then, we can obtain the model existence theorem, which ensures that if we have a tableau with a root formula $@_{i_0}\varphi_0$ one of whose branches is open and saturated, there exists a model which falsifies φ_0 .

Lemma 2 (model existence theorem of TAB_{IB}). Let Θ be an open saturated branch of a tableau in TAB_{IB} and $@_i \varphi$ be a quasi-subformula of the root formula $@_{i_0} \varphi_0$ of Θ where *i* is an identity urfather. Then we have the following proposition:

if $@_i \varphi \in \Theta$, then $\mathcal{M}^{\Theta}, v_{\Theta}(i) \models \varphi$.

Particularly, $\mathcal{M}^{\Theta}, v_{\Theta}(i_0) \models \varphi_0.$

Using this lemma, the completeness theorem for $\mathbf{TAB_{IB}}$ seems to be proved however, an obstacle still remains. The problem we are now facing is though the symmetry of \mathcal{M}^{Θ} is obvious, it is doubtful whether \mathcal{M}^{Θ} is always irreflexive. Unfortunately, the answer is no (consider the model created by the branch in Figure 2.)

To solve this problem, we "bulldoze" all the reflexive points and turn them into an irreflexive frame. For modal logic, the bulldozing method was first used in [19] (see also [14]). For hybrid logic, [4] shows that bulldozing works well in proving the completeness of Hilbert-style axiomatization of irreflexive hybrid logic. Our following lemmata claim that the method also works in tableau calculus.

Definition 15. Given a model $\mathcal{M} = (W, R, V)$ and $W_r = \{w \in W \mid wRw\}$, the bulldozed model $\mathcal{M}_B = (W_B, R_B, V_B)$ is defined as follows:

... For if we take any reflexive world in any model, i.e., any world which can see itself, and replace it with a pair of worlds, each able to see the other but neither able to see itself, and we give each variable the same value in each world in the new pair as it had in the original world, \dots [14, p. 176]

The different point is how we define V(i) where i is nominal: If $V(i) = \{w\}$ and $w \in W_r$, then $V_B(i) = \{(w, 0)\}$.

This construction cannot preserve the satisfaction completely: let i hold in $w \in W$, and w is a reflexive point. Then we have $\mathcal{M}, w \models \Diamond i$, but $\mathcal{M}_B, (w, 0) \models \Diamond i$ does not hold. However, if we consider only the quasi-subformulas of the root formula, this method works well.

First, we divide all the identity urfathers into three groups.

- A) There is another identity urfather j such that $@_i j \in \Theta$.
- B) There is another identity urfather j such that i and j are twins but $T_{\Theta}(i)$ has no nominal.
- C) The others.

Intuitively, each group satisfies the following condition:

- An identity urfather in group A has another identity urfather that points to the same world.
- An identity urfather in group B has another identity urfather such that they point to possibly different but indistinguishable worlds up to quasisubformulas of the root formula.
- Group C is the rest. A member of this group has no identity urfather that points to the same or indistinguishable world.

Using this splitting, we can show the following lemma.

Lemma 3. Let Θ be an open saturated branch, $@_{i_0}\varphi_0$ be a root formula of Θ , and $@_i\varphi$ be a quasi-subformula of $@_{i_0}\varphi_0$ such that $v_{\Theta}(i)$ exists and $v_{\Theta}(i) \in W_r^{\Theta}$. Then, φ cannot be any nominal *i* occurring in φ_0 .

Proof (Sketch). Observe that if $v_{\Theta}(i) \in W_r^{\Theta}$ then it belongs to the Group B (the other cases contradict (\mathcal{I}) .) Since $T_{\Theta}(v_{\Theta}(i))$ has no nominal and so does $T_{\Theta}(i)$, φ cannot be any nominal *i* occurring in φ_0 .

Lemma 4. Let Θ be an open saturated branch, and $@_{i_0}\varphi_0$ be a root formula. For any nominal $i \in W_r^{\Theta}$ and formula φ such that $@_i\varphi$ is a quasi-subformula of $@_{i_0}\varphi_0$, we have

$$\mathcal{M}_B^{\Theta}, (i,0) \models \varphi \quad iff \quad \mathcal{M}_B^{\Theta}, (i,1) \models \varphi.$$

Proof (Sketch). By induction on the complexity of φ . Note that we do not have to consider the case $\varphi = j$ because, by Lemma 3, φ cannot be any nominal *i* occurring in φ_0 .

Lemma 5. Let Θ be an open saturated branch, $@_{i_0}\varphi_0$ be a root formula of Θ , and $@_i\varphi$ be a quasi-subformula of $@_{i_0}\varphi_0$ such that $v_{\Theta}(i)$ exists. Then, we have

$$\mathcal{M}^{\Theta}, v_{\Theta}(i) \models \varphi \text{ implies } \mathcal{M}^{\Theta}_B, (v_{\Theta}(i))_B \models \varphi,$$

where

$$i_B = \begin{cases} \{(i,0)\} & \text{if } i \in W_r, \\ i & \text{otherwise.} \end{cases}$$

Proof (Sketch). We prove it by the induction on the complexity of φ . We give the proof sketch only for the cases $\varphi = \Diamond \psi$.

Suppose that $\mathcal{M}^{\Theta}, v_{\Theta}(i) \models \Diamond \psi$. Then, there is a world $j \in W^{\Theta}$ such that $v_{\Theta}(i)R^{\Theta}j$ and $\mathcal{M}^{\Theta}, j \models \psi$.

The proof is easy if $j \neq v_{\Theta}(i)$. Otherwise, we have $(v_{\Theta}(i), 0)R^{\Theta}(v_{\Theta}(i), 1)$, and $\mathcal{M}_{B}^{\Theta}, (v_{\Theta}(i), 0) \models \psi$ by the induction hypothesis. However, since $@_{v_{\Theta}(i)} \diamond \psi$ is a quasi-subformula of the root formula, so is $@_{v_{\Theta}(i)}\psi$. Thus, by Lemma 4, we have $\mathcal{M}_{B}^{\Theta}, (v_{\Theta}(i), 1) \models \psi$. Therefore, $\mathcal{M}_{B}^{\Theta}, (v_{\Theta}(i), 0) \models \diamond \psi$.

Thus, we have reached our goal.

Theorem 4 (completeness). **TAB**_{IB} is complete for the class of all orthoframes.

Proof. We show the contraposition.

Suppose that φ is not provable in **TAB**_{IB}. Then, we can find a tableau in **TAB**_{IB}, whose root formula is $@_i \neg \varphi$ where *i* does not occur in φ , and which has an open and saturated branch Θ . Then, by Lemma 2, we have $\mathcal{M}^{\Theta}, v_{\Theta}(i) \models \neg \varphi$. By Lemma 5, we have $\mathcal{M}^{\Theta}_B, v_{\Theta}(i)_B \models \neg \varphi (v_{\Theta}(i)_B \text{ is } (v_{\Theta}(i), 0) \text{ if } v_{\Theta}(i) \in W^{\Theta}_r$, and $v_{\Theta}(i)$ otherwise.) Moreover, by the construction, \mathcal{M}^{Θ}_B is an orthomodel. Therefore, we can find an orthoframe which falsifies φ .

Corollary 1. IB(@) is decidable.

Proof. From the termination property of **TAB**_{IB} and the fact that \mathcal{M}_B^{Θ} is finite.

Note that Takeuti and Sano (2020) [16] also shows the decidability of IB(@).

5 Future Work

Besides the satisfaction operator @, hybrid logic can have more operators, such as the existential operator E and the downarrow operator \downarrow . Moreover, hybrid graph logic in [2] contains other modal operators \Box^+ and \diamondsuit^+ , and [16] introduced new modal operators \Box^* and \diamondsuit^* to express the planarity. Since our tableau calculus contains only the basic language, adding more operators is one of our further directions.

Another remaining work is to formulate tableau calculi corresponding to frames with various conditions. Many axioms characterizing some frame conditions are proposed in [8], such as anti-symmetry, trichotomy, and tree-like. If there is a relational structure that is widely studied, creating tableau calculus corresponding to it is worthwhile.

Acknowledgements

We would like to thank Prof. Ryo Kashima and Prof. Katsuhiko Sano for their invaluable advice in writing this paper. The research of the first author was supported by JST SPRING, Grant Number JPMJSP2106. The research of the second author was supported by Grant-in-Aid for JSPS Research Fellow Grant Number JP22KJ1483.

References

- Areces, C., ten Cate, B.: 14 hybrid logics. In: Handbook of Modal Logic, pp. 821– 868. Elsevier (2007)
- Benevides, M.R.F., Schechter, L.M.: Using Modal Logics to Express and Check Global Graph Properties. Logic Journal of the IGPL 17(5), 559–587 (2009)
- Benevides, M.R., Schechter, L.M.: A study on multi-dimensional products of graphs and hybrid logic. Theoretical Computer Science 412, 4946–4966 (2011)
- 4. Blackburn, P.: Nominal tense logic and other sorted intensional frameworks. Ph.D. thesis, The University of Edinburgh (1990)
- Blackburn, P., ten Cate, B.: Pure extensions, proof rules, and hybrid axiomatics. Studia Logica 84(2), 277–322 (2006)
- Blackburn, P., Tzakova, M.: Hybrid languages and temporal logic. Logic journal of IGPL 7(1), 27–54 (1999)
- Bolander, T., Blackburn, P.: Termination for hybrid tableaus. Journal of Logic and Computation 17(3), 517–554 (2007)
- Bolander, T., Blackburn, P.: Terminating tableau calculi for hybrid logics extending K. Electronic Notes in Theoretical Computer Science 231, 21–39 (2009)
- 9. Braüner, T.: Hybrid Logic and its Proof-Theory, vol. 37. Springer Science & Business Media (2011)
- ten Cate, B.: Model Theory for Extended Modal Languages. Ph.D. thesis, University of Amsterdam (2005)
- Gate, J.S.: Applications of Finite Model Theory: Optimisation Problems, Hybrid Modal Logics and Games. Ph.D. thesis, Durham University (2013)

- 12 Y. Nishimura, T. Takagi
- 12. Goldblatt, R.: Logics of Time and Computation. Center for the Study of Language and Information (1987)
- Goldblatt, R.I.: Semantic analysis of orthologic. Journal of Philosophical Logic 3, 19–35 (1974)
- Hughes, G.E., Cresswell, M.J.: A New Introduction to Modal Logic. Psychology Press (1996)
- Indrzejczak, A.: Modal hybrid logic. Logic and Logical Philosophy 16(2-3), 147–257 (2007)
- Izumi, T., Katsuhiko, S.: Modal logic and planarity of graphs. In: Context, Conflict and Reasoning: Proceedings of the Fifth Asian Workshop on Philosophical Logic. pp. 115–126. Springer Singapore (2020)
- 17. Prior, A.N.: Past, Present and Future. Oxford University Press (1967)
- 18. Prior, A.N.: Papers on Time and Tense. Oxford University Press (1968)
- Segerberg, K.K.: An Essay in Classical Modal Logic. Ph.D. thesis, Stanford University (1971)
- Seligman, J., Liu, F., Girard, P.: Logic in the community. In: Proceedings of the 4th Indian Conference on Logic and Its Applications. pp. 178–188. Springer Berlin Heidelberg (2011)
- Seligman, J., Liu, F., Girard, P.: Facebook and the epistemic logic of friendship. In: Proceedings of the 14th Conference on Theoretical Aspects of Rationality and Knowledge. pp. 229–238 (2013)